



TITLE:

# RAAGs in Braids (Complex Analysis and Topology of Discrete Groups and Hyperbolic Spaces)

AUTHOR(S):

KIM, SANG-HYUN; KOBERDA, THOMAS

---

CITATION:

KIM, SANG-HYUN ...[et al]. RAAGs in Braids (Complex Analysis and Topology of Discrete Groups and Hyperbolic Spaces). 数理解析研究所講究録 2015, 1936: 132-136: KJ00009772473.

ISSUE DATE:

2015-04

URL:

<http://hdl.handle.net/2433/223697>

RIGHT:

## RAAGs in Braids

SANG-HYUN KIM\* AND THOMAS KOBERDA

### 1. RIGHT-ANGLED ARTIN GROUPS

In this article, we survey some of the known results regarding right-angled Artin subgroups of right-angled Artin groups and also of mapping class groups. While doing so, we introduce techniques that can improve given embeddings between these groups to simpler ones, which in turn will help us understanding rigidity of embeddings into such groups.

**Definition 1.** Let  $(G, d_G)$  and  $(H, d_H)$  be two groups with metrics. A group homomorphism  $f: G \rightarrow H$  is called a *quasi-isometric group embedding from  $G$  to  $H$*  if  $f$  is injective and there exists  $C \geq 1$  such that for every  $x$  and  $y$  in  $G$  we have

$$d_G(x, y)/C - C \leq d_H(f(x), f(y)) \leq Cd_G(x, y) + C.$$

*Remark.* A finitely generated group will be equipped with a word metric.

For a finite graph  $\Gamma$ , let us define the *RAAG (Right-Angled Artin Group)* on  $\Gamma$  by the group presentation

$$G(\Gamma) = \langle V(\Gamma) \mid [a, b] = 1 \text{ if } \{a, b\} \notin E(\Gamma) \rangle.$$

For example,  $G(\bullet) \cong \mathbb{Z}$ ,  $G(\Delta) \cong F_3$  and  $G(\text{two edges}) \cong F_2 \times F_2$ .

**Question 1.** Which groups arise as subgroups of RAAGs?

**Fact.**

- (1) ([9, 7]) Let  $S$  be a closed surface with  $\xi(S) < -1$ . Then  $\pi_1(S)$  admits a quasi-isometric group embedding into some  $G(\Gamma)$ .
- (2) ([1]) The  $\pi_1$  of every closed hyperbolic 3-manifold virtually admits a quasi-isometric group embedding into some  $G(\Gamma)$ .
- (3) ([3]) For  $d \geq 2$ , there exists a closed hyperbolic  $d$ -manifold  $M_d$  such that  $\pi_1(M_d)$  admits a quasi-isometric group embedding into some RAAG.

**Question 2.**

- (1) Which groups arise as subgroups of a given  $G(\Gamma)$ ?
- (2) Which RAAGs arise as subgroups of a given  $G(\Gamma)$ ?

---

Date: August 20, 2014.

*Key words and phrases.* right-angled Artin group, braid group, cancellation theory, hyperbolic manifold, quasi-isometry.

### Universal property of RAAGs

Suppose  $\phi_1, \dots, \phi_n \in \text{Diff}(M)$  for a manifold  $M$ , and  $\Gamma$  be the intersection graph of  $\{\text{supp}(\phi_1), \dots, \text{supp}(\phi_n)\}$ . Then there exists a “natural” group homomorphism  $G(\Gamma)$  to  $\text{Diff}(M)$ .

**Question 3.** Which RAAGs arise as subgroups of  $\text{Diff}(M)$  or  $\text{Mod}(M)$ ?

## 2. RAAGs IN RAAGs AND IN MODS

*Notation.* For two graphs  $X$  and  $Y$ , we write  $X \leq Y$  if

$$X \subseteq Y \text{ and } EX = EY \cap \binom{VX}{2}.$$

**Theorem 2** ([13]). Let  $\Gamma$  be a finite graph such that  $\mathbb{Z}^3 \not\hookrightarrow G(\Gamma)$ . Then there exists a combinatorially defined, locally infinite graph  $\Gamma^e$  such that for a finite graph  $\Lambda$ , we have

$$G(\Lambda) \hookrightarrow G(\Gamma) \Leftrightarrow \Lambda \leq \Gamma^e.$$

**Theorem 3** ([12] ( $\Rightarrow$ ), [14] ( $\Leftarrow$ )). Let  $S$  be a surface possibly with punctures such that  $\mathbb{Z}^3 \not\hookrightarrow \text{Mod}(S)$ . For a finite graph  $\Lambda$ , we have

$$G(\Lambda) \hookrightarrow \text{Mod}(S) \Leftrightarrow \Lambda^{\text{opp}} \leq C(S).$$

*Remark.* (1) Not true as-is for  $\text{rank} > 2$ . ([6] for RAAGs, [12] for Mods)  
But, there is a version for  $\text{rank} > 2$  using “multi-curves”.

- (2) Abelian ranks are not the only obstructions for  $G(\Gamma) \hookrightarrow \text{Mod}(S)$ .
- (3)  $\Gamma^e$  is a quasi-tree and  $G(\Gamma)$  acts on the opposite graph of  $\Gamma^e$  acylindrically. So we have a “canonical” classification of elements in RAAGs (cf. Bowditch).

## 3. RAAGs ON TREES

**Theorem A** ([11]). For each finite graph  $\Gamma$ , there exists a finite tree  $T$  such that  $G(\Gamma)$  admits a quasi-isometric group embedding into  $G(T)$ .

Here is the recipe. Without loss of generality, we may assume  $\Gamma$  is connected. Consider its universal cover  $p: \tilde{\Gamma} \rightarrow \Gamma$ . For a finite subtree  $T$  of  $\tilde{\Gamma}$ , we have a group homomorphism  $\phi(\Gamma, T): G(\Gamma) \rightarrow G(T)$  defined by

$$\phi(v) = \prod_{u \in p^{-1}(v) \cap T} u.$$

The proof would be complete by showing that for a sufficiently large  $T$ , the map  $\phi(\Gamma, T)$  is a quasi-isometric group embedding.

## 4. APPLICATION I : RAAGS IN BRAIDS

We consider the *pure braid group* on  $n$ -strands:

$$\begin{aligned} PB_n &= \pi_1(\{(z_1, \dots, z_n) \mid z_i \neq z_j\}) \\ &= PMod(D^2 \setminus \{p_1, \dots, p_n\}, \partial D^2). \end{aligned}$$

**Theorem 4** ([8]). *For each finite planar graph  $\Gamma$ , we have a quasi-isometric group embedding from  $G(\Gamma)$  into some pure braid group.*

**Corollary B** ([11]). *Every RAAG admits a quasi-isometric group embedding from into some pure braid group.*

Actually, one can give a self-contained proof.

*proof of Corollary B.* We have only to embed  $G(T)$  for an arbitrary finite tree  $T$ . Consider a collection of disks  $\{D_v \mid v \in V(T)\}$  in  $D^2$  such that the intersection graph is  $T$ . Puncture  $D^2$  sufficiently many times so that there exists a pseudo-Anosov  $\psi_v$  supported on  $D_v \setminus \{\text{punctures}\}$ . There exists a group homomorphism from  $G(T)$  to  $PMod(D^2 \setminus \{\text{punctures}\}) = PB_n$  and this map is a quasi-isometric group embedding, possibly after raising to sufficiently high powers (Clay–Leininger–Mangahas '12).  $\square$

**Question 4** (Farb). *Is the isomorphism problem solvable for f.p. subgroups of  $Mod(S)$ ?*

**Theorem 5** ([4]). (1) *The isomorphism problem is not solvable for f.p. subgroups of a certain  $G(\Gamma_0)$ .*

(2) *The isomorphism problem is not solvable for f.p. subgroups of  $Mod(S_g)$  for  $g \gg 0$ .*

**Corollary 6** ([11]). *The isomorphism problem is not solvable for f.p. subgroups of  $PB_n$  for  $n \gg 0$ .*

## 5. APPLICATION II : SYMPS

We let  $Symp(S^2)$  be the group of area- and orientation-preserving diffeomorphisms (*symplectomorphisms*) of the 2-sphere. For a path  $\{\phi_t\}_{t \in I}$  in  $Symp(S^2)$ , its  $L^p$ -length is defined by

$$\ell_p(\{\phi_t\}) = \int_I \left( \int_{S^2} \left| \frac{\partial \phi_t}{\partial t} \right|^p dx \right)^{1/p}$$

and the (right-invariant)  $L^p$ -metric is given by the corresponding length metric.

**Theorem C** ([11]). *Every RAAG admits a quasi-isometric group embedding into  $(Symp(S^2), d_p)$  for  $p > 2$ .*

*Remark.* (1) ([2, 8]) True for  $D^2$  and  $p \geq 1$ .

(2) Kapovich ('12) showed that every RAAG admits a group embedding into  $Symp(S^2)$  [10].

The proof (independent from Kapovich) goes as follows. From the proof of “RAAGs in braids”, we have a quasi-isometric group embedding

$$\begin{array}{ccc} G(T) & \xrightarrow{\quad} & \mathcal{P}_n \\ & \searrow \text{qi gp emb} & \downarrow q \\ & & \text{PMod}(S^2 \setminus \{p_1, p_2, \dots, p_n\}). \end{array}$$

Here,  $\mathcal{P}_n \leq \text{Symp}(S^2)$  consists of diffeomorphisms that fix some neighborhoods of punctures pointwise. It suffices to show that  $q$  does not “contract too much”. For this, we consider

$$\begin{array}{ccccc} \widetilde{\text{Symp}}(S^2) & \xleftarrow{\text{incl.}} & p^{-1}(\mathcal{P}_n) & \xrightarrow{\quad} & PB_n(S^2) \\ p \downarrow \text{univ. cover, 2-to-1} & & \downarrow \text{2-to-1} & & \downarrow \text{2-to-1} \\ \text{Symp}(S^2) & \xleftarrow{\text{incl.}} & \mathcal{P}_n & \xrightarrow{q} & \text{PMod}(S^2 \setminus \{p_1, \dots, p_n\}) \end{array}$$

The proof would be complete if the upper-right arrow is shown not to contract much. Use Gauss linking integral. This idea is due to (Benaim–Gambaudo ’01, Gambaudo–Ghys ’04, Brandenbursky–Shelukin ’14).

Theorem A is also used in the proof of the following.

**Theorem 7** ([5]). *Every RAAG embeds into  $PL_{\text{Area}}(I^2, \partial I^2)$ .*

Finally, we consider one less dimension:

**Theorem D** (Baik–K.–Koberda). *Every RAAG embeds into  $\text{Diff}^\infty(\mathbb{R})$ .*

**Question 5.** *Does every RAAG embed into  $\text{Diff}^\infty(S^1)$ ?*

Note that every mapping class group embeds into  $\text{Homeo}(S^1)$  but not in  $\text{Diff}^2(S^1)$ . Each RAAG embeds into some mapping class group.

## REFERENCES

1. Ian Agol, *The virtual Haken conjecture*, Doc. Math. **18** (2013), 1045–1087, With an appendix by Agol, Daniel Groves, and Jason Manning. MR 3104553
2. Michel Benaïm and Jean-Marc Gambaudo, *Metric properties of the group of area preserving diffeomorphisms*, Trans. Amer. Math. Soc. **353** (2001), no. 11, 4661–4672 (electronic). MR 1851187 (2002g:58010)
3. Nicolas Bergeron and Daniel T. Wise, *A boundary criterion for cubulation*, Amer. J. Math. **134** (2012), no. 3, 843–859. MR 2931226
4. Martin R. Bridson, *On the subgroups of right angled artin groups and mapping class groups*, To appear in *Math. Res. Lett.*
5. Danny Calegari and Dale Rolfsen, *Groups of PL homeomorphisms of cubes*, preprint (2014).
6. Montserrat Casals-Ruiz, Andrew Duncan, and Ilya Kazachkov, *Embeddings between partially commutative groups: two counterexamples*, J. Algebra **390** (2013), 87–99. MR 3072113
7. John Crisp and Bert Wiest, *Embeddings of graph braid and surface groups in right-angled Artin groups and braid groups*, Algebr. Geom. Topol. **4** (2004), 439–472. MR 2077673 (2005e:20052)

8. ———, *Quasi-isometrically embedded subgroups of braid and diffeomorphism groups*, Trans. Amer. Math. Soc. **359** (2007), no. 11, 5485–5503. MR 2327038 (2008i:20048)
9. Carl Droms, *Graph groups, coherence, and three-manifolds*, J. Algebra **106** (1987), no. 2, 484–489. MR 880971 (88e:57003)
10. Michael Kapovich, *RAAGs in Ham*, Geom. Funct. Anal. **22** (2012), no. 3, 733–755. MR 2972607
11. Sang-hyun Kim and Thomas Koberda, *Anti-trees and right-angled Artin subgroups of planar braid groups*, preprint.
12. ———, *Right-angled Artin groups and finite subgraphs of curve graphs*, preprint.
13. ———, *Embedability between right-angled Artin groups*, Geom. Topol. **17** (2013), no. 1, 493–530. MR 3039768
14. Thomas Koberda, *Right-angled Artin groups and a generalized isomorphism problem for finitely generated subgroups of mapping class groups*, Geom. Funct. Anal. **22** (2012), no. 6, 1541–1590. MR 3000498

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, DAEJEON 305-701, REPUBLIC OF KOREA

*E-mail address:* `shkim@kaist.edu`

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, 20 HILLHOUSE AVE, NEW HAVEN, CT 06520, USA

*E-mail address:* `thomas.koberda@gmail.com`